

BREAKDOWN OF ZERO-MODE APPROXIMATION IN THE INSTANTON VACUUM

B.O.Kerbikov, D.S.Kuzmenko, Yu.A.Simonov
Institute of Theoretical and Experimental Physics,
117259 Moscow, Russia

A new approach to effective theory of quarks in the instanton vacuum is presented. Exact equations for the quark propagator and Lagrangian are derived which contain contributions of all quark modes with known coefficients. The resulting effective Lagrangian resembles that of the Nambu–Jona–Lasinio model. The commonly used zero-mode approximation is shown to be invalid in the chiral limit.

PACS: 02.30.Jr, 11.15.-q

1. The recent lattice data [1] provided evidence that instantons may be responsible for nonperturbative behavior of $q\bar{q}$ correlators [2] which makes the study of the quark dynamics in the instantonic vacuum [3–5] an important and fundamental problem.

To date practically all papers on the subject have relied upon the use of the so-called zero-mode approximation (ZMA) which amounts to comprising only zero quark mode in a single-instanton fermion propagator [3]. Correspondingly an ansatz for the partition function and effective quark Lagrangian (EQL) containing zero modes only have been proposed [5] and widely used in literature [6,7].

The purpose of this letter is to give a complete normal mode expansion of the EQL and of the quark propagator. For a specific choice of the coefficients in this expansion one retrieves the ansatz [5] for EQL. However this option is invalidated by exact calculation presented below which shows that zero-mode term vanishes in the chiral limit and hence ZMA for EQL breaks down. More intricate is the analysis of the quark propagator S in the instanton–anti-instanton vacuum. Here the zero-mode term survives but enters with a coefficient depending solely on contributions of higher modes. The same feature can be seen in the quark partition function $\langle \det S^{-1} \rangle$, where the average $\langle \dots \rangle$ is defined below. Thus ZMA breaks down and the new quark dynamics associated with nonzero modes emerges. The main features of this dynamics are outlined in what follows.

2. To make discussion transparent consider an ideal instanton gas with the superposition ansatz [8–10] and with equal number of instantons and anti-instantons, $N_+ = N_- = N/2$:

$$A_\mu(x) = \sum_{i=1}^N A_\mu^{(i)}(x - R_i), \quad (1)$$

$$gA_\mu^{(i)} = \frac{\bar{\eta}_{a\mu\nu}(x - R_i)_\nu \rho^2 \Omega_i^+ \tau_a \Omega_i}{(x - R_i)^2 [(x - R_i)^2 + \rho^2]}, \quad (2)$$

where Ω_i , R_i , and ρ are color orientation, position and scale-size of the i -th instanton.

The EQL is obtained from the Euclidean partition function after averaging over $\{\Omega_i, R_i\}$:

$$Z = \int D\psi D\psi^+ e^{-\int dx \psi^+ S^{-1} \psi} \prod_{i=1}^N \frac{dR_i}{V} d\Omega_i = \int D\psi D\psi^+ e^{-L_{eff}}, \quad (3)$$

where definitions here and in what follows are:

$$\begin{aligned} S_0^{-1} &= (-i\hat{\partial} - im_f), \\ S_i^{-1} &= (-i\hat{\partial} - g\hat{A}^{(i)} - im_f), \\ S^{-1} &= (-i\hat{\partial} - g\hat{A} - im_f). \end{aligned} \quad (4)$$

Next we introduce the standard set of eigenfunctions $\{u_n^i\}$, $n = 0, 1, 2, \dots$

$$(-i\hat{\partial} - g\hat{A}^{(i)})|u_n^i\rangle = \lambda_n|u_n^i\rangle, \quad (5)$$

and expand S^{-1} given by (4) as

$$S^{-1} = S_0^{-1} + \sum_{i,m,n} S_0^{-1}|u_m^i\rangle \varepsilon_{mn}^i \langle u_n^i| S_0^{-1}, \quad (6)$$

where $\hat{\varepsilon}$ can be represented either as

$$\varepsilon_{mn}^i = \langle u_m^i | S_0 S_i^{-1} S_0 - S_0 | u_n^i \rangle = - \langle u_m^i | (S_i - S_0) [1 + S_0^{-1} (S_i - S_0)]^{-1} | u_n^i \rangle, \quad (7)$$

or simply as

$$\varepsilon_{mn}^i = -g \langle u_m^i | S_0 \hat{A}^{(i)} S_0 | u_n^i \rangle. \quad (8)$$

Performing in (3) averaging with the help of cumulant or cluster expansion, one obtains L_{eff} in the form

$$L_{eff} = - \int dx \psi^+ S_0^{-1} \psi + \sum_{n=2}^{\infty} (-1)^n 2^n \left(\frac{V}{N}\right)^{n-1} \sum_{fmm'} \int d\Gamma_n \det_{k,l}^{(n)} J_{kl}, \quad (9)$$

where

$$d\Gamma_n = \prod_{j=1}^n \frac{dp_j}{(2\pi)^4} \cdot \frac{dp'_j}{(2\pi)^4} (2\pi)^4 \delta \left(\sum_j (p_j - p'_j) \right), \quad (10)$$

$$J_{kl} = \left(\psi^{fk}(p_k) \right)^+ M_{mm'}^{fk f_l}(p_k, p'_l) \psi^{f_l}(p'_l), \quad (11)$$

and similarly to [5,6] we have introduced the vertices

$$M_{mm'}^{gr}(p, p') = \frac{N}{2VN_c} (\hat{p} - im_g) \varphi_m(p) \varepsilon_{mm'}^i \varphi_{m'}^+(p') (\hat{p}' - im_r), \quad (12)$$

with $\varphi_m(p)$ being the form factor of u_m^i in momentum space.

Summation in (9) starts from $n = 2$ since the $n = 1$ term drops out as a result of integration over color orientations.

The EQL in (9) is a sum of $n \times n$ determinants. If one confines oneself to ZMA i.e. puts ε_{00}^i finite and $\varepsilon_{m>0,n>0}^i$ equal to zero the sum runs only over $n \leq N_f$. This restriction is due to Grassmann nature of J_{kl} . Thus even in ZMA one obtains e.g. for $N_f = 3$ three 2×2 determinants and one 3×3 determinant. Only the last one is present in the ansatz [5] with the identification $\varepsilon_{00}^i \equiv \varepsilon$, $M_{00} \equiv M$. Therefore our results are in contrast to the common lore according to which for a given number of flavors N_f the only vertex appearing in the chiral limit contains $2N_f$ quark operators. We can reproduce this result for $N_f = 2$ if only ε_{00}^i is kept nonzero, while for $N_f = 3$ this conjecture does not suffice and we get additional $4q$ terms.

Now we come to the central point of the letter. Let us estimate ε_{00}^i using (8). In the chiral limit the term $S_0 \hat{A}^{(i)} S_0$ is chirally odd and therefore ε_{00}^i vanishes, for $m_f \neq 0$ one has

$$\varepsilon_{00}^i = O(m_f), \quad m_f \rightarrow 0. \quad (13)$$

At the same time nonzero modes u_{mn}^i do not have definite chirality and hence matrix elements ε_{mn}^i do not vanish as $m_f \rightarrow 0$. Thus all terms proportional to ε_{00}^i in L_{eff} vanish in the chiral limit implying the breakdown of ZMA in the effective quark Lagrangian.

3. Now we turn to the quark propagator, expressing it again through ε_{mn}^i . Inverting (6) one finds

$$S = S_0 - \sum_{ijmn} |u_m^i\rangle \left(\frac{1}{\hat{\varepsilon}^{-1} + \hat{V}} \right)_{mn}^{ij} \langle u_n^j|, \quad (14)$$

where $(\hat{\varepsilon})_{mn}^{ij} = \delta_{ij} \varepsilon_{mn}^i$, and

$$(\hat{V})_{mn}^{ij} = \langle u_m^i | S_0^{-1} | u_n^j \rangle. \quad (15)$$

Note that summation in (14) extends over different instantons and hence over u_0^i and u_0^j of different chiralities. Equation (14) has to be compared to the following expression common to most papers on the subject [3,5,6]

$$S = S_0 - \sum_{i,j} |u_0^i\rangle \left(\frac{1}{2im + V} \right)_{00}^{ij} \langle u_0^j|, \quad (16)$$

which contains only zero modes contributions. To derive (16) one starts with the following approximation for the quark propagator in a single instanton field [3,5]:

$$S_i = (-i\hat{\partial})^{-1} + \frac{|u_0^i\rangle \langle u_0^i|}{-im}. \quad (17)$$

Introducing this ansatz into expression (7) for ε_{mn}^i we get

$$\varepsilon_{00}^i = \frac{1}{2im}, \quad \varepsilon_{m>0,n>0}^i = 0. \quad (18)$$

Using this form of $\hat{\varepsilon}$ in (14) one recovers the standard ZMA (16). Now, comparing (18) to (13) we conclude that ansatz (17) is unjustified. Actually when ε_{00}^i vanishes in the chiral limit in line

with (13), the propagator (14) still contains terms $|u_0^i \rangle \langle u_0^j|$, but with coefficients depending upon higher modes contributions V_{mn}^{ij} . This can be seen expanding (14) in series in powers of ε , i.e.

$$S = S_0 - \sum_{i,j,m,n} |u_m^i \rangle (\hat{\varepsilon} - \hat{\varepsilon}\hat{V}\hat{\varepsilon} + \hat{\varepsilon}\hat{V}\hat{\varepsilon}\hat{V}\hat{\varepsilon} - \dots)_{mn}^{ij} \langle u_n^j|. \quad (19)$$

If one neglects nonzero modes in V_{mn} in (21), then the coefficient of $|u_0^i \rangle \langle u_0^j|$ automatically vanishes. Thus the popular ZMA for the propagator (16), using only zero modes for different instantons and anti-instantons, loses its ground—this contribution vanishes in the chiral limit while nonzero modes give a nonvanishing result.

It is worth noting that the consistency of the approximation (17) was questioned in [9] in connection to the calculation of the two-point correlation function. It was shown in [9] that it is absolutely necessary to keep the order $\sim m$ terms in S_i . However since for massive fermions the single instanton propagator S_i is not explicitly known the effects of higher modes and finite mass have been investigated only numerically [11].

Finally let us see the effect of nonzero modes in the quark partition function, which is obtained from (3) integrating first over quark fields. Using (6) for S^{-1} one easily obtains

$$Z/Z_0 = \prod_{f=1}^{N_f} \det(1 + \hat{\varepsilon}\hat{V}), \quad (20)$$

where $\hat{\varepsilon}$ and $\hat{V}$ are the same matrices as in (14), (15). It is clearly seen in (20) that the contribution of only zero modes on different instantons, i.e. V_{00}^{ij} is accompanied by factors ε_{00}^k , and vanishes in the chiral limit due to (13). Thus in this limit nonzero modes are necessary to ensure nonvanishing result.

4. One may wonder, why ZMA is invalid even though phenomenologically it looks like giving reasonable results [5-6, 12]. One of the reasons might be that $\varepsilon_{00}^i \equiv \varepsilon$ has been treated as a parameter connected to the properties of the instanton vacuum via the relation $\varepsilon \sim (\frac{N_c V}{N \rho^2})^{1/2}$, while the properties of the vacuum have been in turn adjusted to the correct value of the gluon condensate.

Our results are at first sight in contradiction to the Banks–Casher relation [13] which connects the chiral condensate with the density of global (quasi) zero modes. The standard picture suggests that the later originate from individual zero modes, and hence would disappear as soon as (13) holds. However here the standard picture fails. An insight into its failure is provided by quantum mechanics of collective levels in N potential wells in $4d$. If each of the well has one loosely bound level and continuum (equivalent to zero mode and nonzero modes), then the approximation of keeping only the bound state poles in the Green's functions of each well is known to give an inadequate description of collective bound states [14]. More than that, the pole approximation is a poor one even for the Green's function of the individual well, and instead the so-called unitary pole approximation (UPA) has to be used [15].

5. To summarize, we have outlined the new approach to the effective theory of quarks in the instanton vacuum. Our EQL is similar to that of NJL [16], namely it starts from $4q$ term which might play an important role in phenomenology. Analogy to NJL model calls for construction

of gap equation yielding chiral quark mass and quark condensate. Finally, the bosonization procedure has to be performed yielding the effective chiral Lagrangian for Nambu–Goldstone modes. This program is in progress now and will be reported elsewhere.

The authors are thankful to A.V.Smilga, Yu.M.Makeenko and S.V.Bashinsky for useful discussions. The work was supported by INTAS Grant 94–2851 and by the Russian Fund for Fundamental Research Grant 96–02–19184a.

-
1. M.C.Chu, J.M.Grandy, S.Huang, J.W.Negele, Phys. Rev. Lett. **70**, 225 (1993); Phys. Rev. **D49**, 6039 (1994).
 2. E.V.Shuryak, Rev. Mod. Phys. **65**, 1 (1993).
 3. D.Diakonov and V.Petrov, Nucl. Phys. **B272**, 457 (1986).
 4. E.V.Shuryak, Nucl. Phys. **B302**, 559, 574, 599 (1988).
 5. D.Diakonov and V.Petrov, Preprint LNPI–1153 (1986);
D.Diakonov and V.Petrov, in: Quark Cluster Dynamics, Lecture Notes in Physics, Springer–Verlag (1992) p.288.
 6. M.A.Nowak, J.J.M.Verbaarschot, and I.Zahed, Phys. Lett. **B228**, 251 (1989); Nucl. Phys. **B324**, 1 (1989).
 7. Yu.A.Simonov, Yad. Fiz. **57**, 1491 (1994).
 8. C.Callan, R.Dashen, and D.Gross, Phys. Rev. **D17**, 2717 (1978).
 9. N.Andrei and D.J.Gross, Phys. Rev. **D18**, 468 (1978).
 10. H.Levin and L.G.Yaffe, Phys. Rev. **D19**, 1225 (1979).
 11. E.V.Shuryak and J.J.M.Verbaarschot, Nucl. Phys. **B410**, 37 (1993).
 12. T.Wettig, A.Schäfer, and H.A.Weidenmüller, Phys. Lett. **B367**, 28 (1996).
 13. T.Banks and A.Casher, Nucl. Phys. **B169**, 103 (1980).
 14. A.I.Baz', Ya.B.Zeldovich, and A.M.Perelomov, *Scattering, reactions and decays in non-relativistic quantum mechanics*, Moscow, Nauka, 1971;
B.Sakita, *Quantum theory of many-variable systems and fields*, World Scientific, 1985.
 15. C.Lovelace, Phys. Rev. **135B**, 122 (1964).
 16. Y.Nambu and G.Jona–Lasinio, Phys. Rev. **122**, 345 (1961).